

# On the bootstrap structure of Yangian-invariant factorized S-matrices

Niall J. MacKay\*

*Department of Mathematical Sciences, Durham, DH1 3LE, England*

The message of this talk is that there is some very rich, largely unremarked structure in the representation theory of the Yangian (a quasitriangular Hopf algebra discovered by Drinfeld<sup>[1]</sup> in 1985).

We begin by taking the point of view<sup>[2,3]</sup> of a physicist investigating the charge algebra of integrable field theories in 1+1 dimensions which have a Lie algebra valued, curvature-free, conserved current (of which notable examples are the principal chiral model and the Gross-Neveu model and its generalizations). Such theories thus contain a charge  $Q_0^a$ , valued in a Lie algebra  $\mathcal{A}$ , which acts additively on tensor products of asymptotically independent particle states: defining its action on a two-particle state to be  $\Delta(Q_0^a)$ , we have

$$[Q_0^a, Q_0^b] = i\hbar f^{abc} Q_0^c \quad \text{and} \quad \Delta(Q_0^a) = Q_0^a \otimes 1 + 1 \otimes Q_0^a . \quad (1)$$

Such theories also have higher *non-local* conserved charges. The first of these,  $Q_1^a$ , satisfies

$$[Q_0^a, Q_1^b] = i\hbar f^{abc} Q_1^c \quad \text{and} \quad \Delta(Q_1^a) = Q_1^a \otimes 1 + 1 \otimes Q_1^a + \frac{1}{2} f^{abc} Q_0^c \otimes Q_0^b . \quad (2)$$

The missing commutator of two  $Q_1$ 's is fixed by requiring that asymptotic states carry a representation of the charge algebra (*i.e.* that  $\Delta$  be a homomorphism), and is ( $\mathcal{A} \neq sl(2)$ )

$$f^{d[ab}[Q_1^c], Q_1^d] = \frac{i\hbar}{12} f^{api} f^{bjq} f^{crk} f^{ijk} Q_0^p Q_0^q Q_0^r . \quad (3)$$

This complicated relation may safely be ignored for the moment, although we shall return to it later. The Lorentz boost does not act trivially on these charges. If we boost the rapidity (defined by  $\mathbf{p} = (m \cosh\theta, m \sinh\theta)$ ) of a particle by  $\theta$ , it is found that

$$L_\theta : Q^{(0)} \mapsto Q^{(0)} , \quad Q^{(1)} \mapsto Q^{(1)} - \frac{\hbar C_{Adj}}{4\pi} \theta Q^{(0)} \quad (\text{where } C_{Adj} \delta_{ad} = f^{abc} f^{abd}) . \quad (4)$$

This has important implications for the  $S$ -matrix, which is heavily constrained by conservation of  $Q_0$  and  $Q_1$ . The  $S$ -matrix for the interaction of two particles of rapidities  $\theta_1$  and  $\theta_2$  must satisfy

$$[S(\theta_1 - \theta_2), Q_0^a \otimes 1 + 1 \otimes Q_0^a] = 0$$

$$S(\theta_1 - \theta_2) (L_{\theta_1} \otimes L_{\theta_2} \Delta(Q_1^a)) = (L_{\theta_2} \otimes L_{\theta_1} \Delta(Q_1^a)) S(\theta_1 - \theta_2) . \quad (5)$$

---

\*Supported by a UK SERC studentship.


These equations, together with unitarity and crossing-symmetry, determine the  $S$ -matrix up to an overall scalar factor. The classic result implied by these equations is that the  $S$ -matrix is *factorized*: that the  $S$ -matrix for the interaction of a large number of particles may be written as a product of two-particle  $S$ -matrices. That this works consistently follows from the Yang-Baxter equation (YBE),

$$S_{V_2 V_3}(\theta') S_{V_1 V_3}(\theta + \theta') S_{V_1 V_2}(\theta) = S_{V_1 V_2}(\theta) S_{V_1 V_3}(\theta + \theta') S_{V_2 V_3}(\theta') \quad (6)$$

(where  $\theta$  and  $\theta'$  denote the appropriate rapidity differences), which follows from (5): the  $S$ -matrix does not depend on the order in which the particles interact, and its only action is to exchange quantum numbers within the multiplets  $V_i$  of particles of equal mass  $m_i$ . Factorization was actually deduced some years ago from other considerations, and such  $S$ -matrices have been studied for some time<sup>[4,5]</sup>. In fact, as Bernard<sup>[3]</sup> pointed out, the algebra (1,2,3) is precisely Drinfeld's Yangian  $Y(\mathcal{A})$ , which is a quasitriangular Hopf algebra with an  $R$ -matrix satisfying (5,6) (where  $S = \mathbf{P}R$  with  $\mathbf{P}$  the transposition map), and has a one-parameter automorphism (4). Now suppose that we wish to construct  $S$ -matrices without using the underlying Yangian structure. In order to do so we may use the *bootstrap principle*: that, at appropriate poles, intermediate states of the  $S$ -matrix should be regarded as physical particles. For example, where  $\mathbf{1}$  is the particle in the vector representation of  $SO(N)$  we have<sup>[4]</sup>

$$S_{\mathbf{1}\mathbf{1}}(\theta) = s(\theta) [(\theta_0 - \theta)(i\pi - \theta)P_S + (\theta_0 + \theta)(i\pi - \theta)P_A + (\theta_0 + \theta)(i\pi - \theta)P_0]$$

where  $P_S, P_A$  and  $P_0$  project onto the traceless symmetric, antisymmetric and trace components of  $\mathbf{1} \otimes \mathbf{1}$ ,  $\theta_0 = \frac{2i\pi}{N-2}$ , and  $s(\theta)$  is a scalar function. The only value of  $\theta$  in the physical strip  $0 \leq \text{Im}\theta < \pi$  at which any of these components vanish is  $\theta = \theta_0$ , and we therefore choose the ambiguous factor  $s(\theta)$  to have a simple pole at  $\theta_0$ , and interpret  $P_A + P_0$  as a particle state  $\mathbf{2}$ , so that we have a 'fusing'  $\mathbf{1}\mathbf{1} \rightarrow \mathbf{2}$ . Conservation of momentum tells us the relative masses of these particles, and we can construct  $S_{\mathbf{1}\mathbf{2}}$  and  $S_{\mathbf{2}\mathbf{2}}$  according to the principle<sup>[6]</sup>



We can now continue the bootstrap, examining  $S_{\mathbf{1}\mathbf{2}}$  and  $S_{\mathbf{2}\mathbf{2}}$  for poles, interpreting these as third and higher particles, and so on. In principle, we would hope that eventually all

appropriate poles would have been interpreted as physical states, so that the bootstrap would have closed on a finite set of particles. The mass ratios of the particles would be known, as would the representation (possibly reducible) of  $\mathcal{A}$  carried by the particle, and the set of particle ‘fusings’  $\mathbf{ab} \rightarrow \mathbf{c}$ . Unfortunately, calculating the  $S$ -matrices involves a great deal of computation, and calculations beyond  $S_{\mathbf{22}}$  are unfeasible in practice. However, we can already extract one important point. In all cases hitherto computed, it has been possible to assign a particle  $\mathbf{i}$  to a reducible representation of  $\mathcal{A}$  containing the  $i$ th fundamental representation  $V_i$  of  $\mathcal{A}$ . In some cases (*e.g.* for  $\mathcal{A} = a_n$  and for vector and spinor representations of  $SO(N)$ ) there are no other components. A general method exists for calculating the  $S$ -matrices in irreducible representations<sup>[7]</sup>, and the bulk of the  $S$ -matrices so far constructed have been of this type<sup>[5,7]</sup>, and led to the conjecture that the fusings are in precise agreement with the Clebsch-Gordan (CG) decomposition, *i.e.* that  $\mathbf{ab} \rightarrow \mathbf{c}$  if and only if  $V_a \otimes V_b \supset V_c$ . However, when the particle multiplet is reducible, this fails: for example, when we construct  $S_{\mathbf{22}}$  for the  $SO(N)$  theories<sup>[8]</sup>, we find that  $\mathbf{22} \not\rightarrow \mathbf{2}$  in general even though  $V_2 \otimes V_2 \supset V_2$ .

Now let us compare these  $S$ -matrices with another type of factorized  $S$ -matrix. In *purely elastic scattering theories* or PESTs (of which the prime examples are affine Toda theories<sup>[9]</sup> and integrable deformations of CFTs<sup>[10]</sup>), there is no mass-degeneracy: the particles do not form multiplets, and (6) is trivial. The  $S$ -matrix is a scalar factor determined by unitarity, analyticity, crossing-symmetry and the bootstrap, which is now much easier to implement. The result is a closed bootstrap on a full spectrum of massive particles and their fusings. For simply-laced  $\mathcal{A}$  these match precisely the known results for the YBE-dependent  $S$ -matrices (including the CG ‘holes’)<sup>†</sup>; moreover, the PEST structure has a beautiful description in terms of root systems of Lie algebras<sup>[11]</sup>. It therefore seems likely that similar structure exists in the YBE-dependent  $S$ -matrices - it is simply that the bootstrap, whose implementation is specific to the  $\mathcal{A}$  under consideration, is unable to perceive it.

Let us therefore shift viewpoint and consider the particle multiplets  $\mathbf{i}$  as fundamental representations of  $Y(\mathcal{A})$ . We can use (4) to boost  $\mathbf{i} \mapsto \mathbf{i}(\theta)$ ; the fusion procedure for  $S_{\mathbf{ab}}(\theta)$  is then that of decomposing  $\mathbf{a}(\theta) \otimes \mathbf{b}$  into its  $Y(\mathcal{A})$ -irreducible components. An alternative approach to constructing  $S$ -matrices is to construct the action of  $Q_0$  and  $Q_1$  explicitly and solve (5). This has been done, as far as I am aware, for two cases only. The first is when  $\mathbf{i} = V_i$  is irreducible and  $\rho_{\mathbf{i}}(Q_1) = 0$ : this only works when the action of the right-hand side of (3) is zero. The second is the extension of the adjoint representation of  $\mathcal{A}$  by a scalar representation to the  $Y(\mathcal{A})$ -irreducible representation  $\mathcal{A} \oplus \mathbf{C}$ : the action

---

<sup>†</sup>For non-simply-laced  $\mathcal{A}$  there does seem to be a relationship, but it is much less clear.

of  $Q_0$  is the adjoint action on  $\mathcal{A}$  and zero on  $\mathbf{C}$ , and the action of  $Q_1$  is then constructed to satisfy (2,3). Both of these types of representation were discovered by Drinfeld<sup>[1]</sup>, and the corresponding  $R$ -matrices have been calculated<sup>[7,12]</sup>. Unfortunately this is where the story currently ends. The construction of other representations of  $Y(\mathcal{A})$  seems to me to be an interesting problem for mathematicians: for example, can we construct  $\rho_i$  by taking  $Q_0$  to have the natural action, and then fixing the action of  $Q_1$  so that (2,3) are satisfied? Beyond this, solving (5) for  $S$  is certain to be a tricky process, and may yield little insight into the information obtained; at all levels, it seems, generality is lacking.

In conclusion, there is every prospect of interesting structure in the representation theory of  $Y(\mathcal{A})$  alongside a dearth of general techniques for its discovery. For mathematical physicists, I should perhaps point out in addition the wide range of 1+1 dimensional integrable theories to which  $Y(\mathcal{A})$  is relevant - apart from those mentioned at the beginning, the close relationship with PESTs has even led Belavin<sup>[13]</sup> to suggest that the underlying symmetry of affine Toda theories may be  $Y(\mathcal{A})/\mathcal{A}$ . I can only echo this, and hope to have convinced people that the subject deserves more attention than it has hitherto received.

## References

- [1] V. G. Drinfeld, Sov.Math.Dokl. 32 (1985) 254.
- [2] M. Lüscher, Nucl.Phys. B135 (1978) 1.
- [3] D. Bernard, Comm.Math.Phys. 137 (1991) 191.
- [4] A. B. Zamolodchikov and A. B. Zamolodchikov, Ann.Phys. 120 (1979) 253.
- [5] E. Ogievetsky and P. Wiegmann, Phys.Lett. B168 (1986) 360.
- [6] M. Karowski, Nucl.Phys. B153 (1979) 244.
- [7] P. P. Kulish, E. K. Sklyanin and N. Y. Reshetikhin, Lett.Math.Phys. 5 (1981) 393.
- [8] N. J. MacKay, Nucl.Phys. B356 (1991) 729.
- [9] H. W. Braden, E. Corrigan, P. E. Dorey and R. Sasaki, Nucl.Phys. B338 (1990) 689.
- [10] A. B. Zamolodchikov, Int.J.Mod.Phys A4 (1989) 4235.
- [11] P. E. Dorey, Nucl.Phys. B358 (1991) 654.
- [12] V. Chari and A. Pressley, J.für die reine und ang. Math. 417 (1991) 87.
- [13] A. A. Belavin, Phys.Lett. 283B (1992) 67.